

References

1. Birch C. S., Bonwick G. A. // International Journal of Food Science & Technology. 2018. P. 1467–1485.
2. Albert C., Gombos S., Salamon R. V. et al. // Acta Universitatis Sapientiae, Alimentaria. 2017. P. 5–20.
3. Velázquez-Contreras F., Acevedo-Parra H., Nuño-Donlucas S. M. et al. // Polymers. 2019. Vol. 11. P. 1720–1730.
4. Shi M., Jing Y., Yang L. et al. // Polymers. 2019. Vol. 11. P. 14.
5. Nara S., Mori A., Komiya T. // Starch. 1978. Vol. 4. P. 111–114.

* This research was carried out within the State Assignment to ISSCM SB RAS (project 0301-2019-0004).

УДК 548.73:547.221'442.3'546.34

V. I. Filyakova


*I. Ya. Postovsky Institute of Organic Synthesis,
Ural Branch of the Russian Academy of Sciences,
620108, Russia, Ekaterinburg,
S. Kovalevskoy / Akademicheskaya St., 22 / 20,
vif@ios.uran.ru*

LITHIUM 3-POLYFLUOROALKYL-1,3-DIKETONATES AS A VALUABLE BUILDING BLOCKS*


Key words: Lithium 3-polyfluoroalkyl-1,3-diketonates, 4-amino-pyrazoles, quinoxaline and pyridine derivatives, terpyridine.

Previously, we have introduced lithium 3-polyfluoroalkyl-1,3-diketonates (LDKs) in organic synthesis as building blocks and demonstrated benefits of their use compared to appropriate 1,3-diketones. They are readily accessible, stable on storage, and highly reactive. On the bases on LDKs, we have developed facile syntheses methods of fluoroalkyl-containing polyfunctional compounds [1–8], various linear and annulated heterocycles [1, 9–14].

We have significantly expanded the synthetic capabilities of LDKs **1** through nitrosation reactions and developed effective methods for the synthesis of 5-hydroxy-5-(polyfluoroalkyl)isoxazol-4(5*H*)-one oximes **3** [14], 4-amino-pyrazoles **4**, quinoxaline derivatives **5** and **6** [14] and new heterocyclic systems **7** and **8**.



In addition, when LDKs were reacted with ammonium acetate in AcOH, pyridine derivatives **9** were unexpectedly obtained. Of particular importance is the process of converting LDK **1b** to terpyridine **9b**, which is an effective ligand for the design supramolecular systems, creating organic chemosensors, photosensitizers, etc.



References

1. Filyakova V. I., Karpenko N. S., Kuznetsova O. A., Pashkevich K. I. // *Russ. J. Org. Chem.* 1998. Vol. 34. P. 381.
2. RU Patent 2100345, Byul. Izobr. 36 (1997) (in Russian).
3. Boltacheva N. S., Filyakova V. I., Charushin V. N. // *Russ. J. Org. Chem.* 2005. Vol. 41, № 3. P. 1452.
4. Boltacheva N. S., Fedorova O. V., Ovchinnikova I. G. et al. // *J. Fluor. Chem.* 2007. Vol. 128. P. 762.
5. Filyakova V. I., Boltacheva N. S., Pervova M. G. et al. // *Mend. Comm.* 2017. Vol. 27. P. 464.
6. Filyakova V. I., Kuznetsova O. A., Ulomsky E. N. et al. // *Russ. Chem. Bull.* 2002. Vol. 51. P. 332.
7. Sevenard D. V., Khomutov O. G., Boltachova N. S., Filyakova V. I. et al. // *Z. Naturforsch.* 2009. Vol. 64b. P. 541.
8. Filyakova V. I., Boltacheva N. S., Sevenard D. V., Charushin V. N. // *Russ. Chem. Bull.* 2010. Vol. 59, № 9. P. 1791.
9. Boltacheva N. S., Dianova L. N., Slepukhin P. A., Filyakova V. I., Charushin V. N. // *Russ. J. Org. Chem.* 2014. Vol. 50, № 6. P. 785.
10. Palysaeva N. V., Boltacheva N. S., Slepukhin P. A., Pervova M. G., Filyakova V. I., Sheremetev A. B., Charushin V. N. // *Mend. Comm.* 2018. Vol. 28. P. 126.

11. Boltacheva N. S., Filyakova V. I., Charushin V. N. // RU Patent № 2642924. 2018. // Bull. Izobret. 2018. № 4.
12. Filyakova V. I., Boltacheva N. S., Charushin V. N. // Russ. J. Org. Chem. 2019. Vol. 55, № 6. P. 890–893.

** This work was supported by a grant RFBR (Project № 18-03-00 112 A).*

УДК 547.7


**M. N. Joy, N. A. Beliaev,
T. V. Beryozkina, V. A. Bakulev**

*TOS Department, Ural Federal University,
620002, Russia, Yekaterinburg, Mira St., 19,
v.bakulev@urfu.ru; mnibinjoy@urfu.ru*

FACILE ONE-POT SYNTHESIS OF PHARMACOLOGICALLY RELEVANT 1,2,3-TRIAZOLES LINKED TO EQUOL*

Keywords: equol, cycloaddition, 1,2,3-triazole, copper.

A convenient access for the synthesis of some pharmacologically relevant equol derivatives linked to 1,2,3-triazoles has been achieved [1]. The salient features of this developed protocol include: easy isolation process, good to excellent yield of the products and appendage diversity of heteroaryl triazoles (scheme).



References

1. Joy M. N., Bodke Y. D., Telkar S., Bakulev V. A. // Journal of the Mexican Chemical Society. 2020. Vol. 64. P. 46–66.

** The work was supported by the RFBR grant № 170300641A.*